


MHC Class II Capture Kit

Catalog No.	32015S	32015L
Reactions	24 reactions	96 reactions

Description

The **MHC Class II Capture Kit** was developed to capture the entire MHC Class II locus from whole genome NGS libraries based on the CATCH-Seq technology. Comprehensive sequencing of the entire Class II locus with our kit make it possible to detect SNPs, indels, and structural variants that are not covered by other MHC targeted sequencing reagents.



Our MHC Class II kit enables the most cost-effective targeted sequencing of 0.7 Mb of the human MHC Class II locus, including all coding and non-coding regions.

Captured sample multiplexing:

- **Single Index** (Cat. # 30072 Multiplexing Index Primers): For kits with 24 reactions.
- **Unique dual index** (Cat. # 30075 Multiplexing Unique Dual Index Primers): For kits with 96 reactions.

Features

- **Full MHC Class II region**
 - Covers 0.7 MB of MHC Class II region
 - Covers exons, introns, 5' regulatory regions, 3' regulatory regions, and beyond.
- **Easy detection of SNPs, indels, and structural variants**
 - The only reagent provides intact sequence information
- **Low cost**
- **Sample multiplexing:** Further reduces the cost

Component

Catalog No.	32015S	32015L
MHC-II-B Probe	1440 ul	5760 ul
RS Buffer	672 ul	2688 ul
W1 Buffer	14.4 ml	57.6 ml
W2 Buffer	9.6 ml	38.4 ml

Storage Condition

- Store MHC-II-B Probe at -20°C, stable up to 12 months.
- Store W1, W2 and RS Buffer at room temperature, stable up to 12 months.

Reagent & Equipment Needed (not provided in this kit)

- Magnetic particle concentrator
- PCR thermal cycler
- Hybridization oven
- Heat block (compatible with 96-well plate)
- 96-well PCR plate
- 80% ethanol (prepare before use)
- MyOne streptavidin C1 dynabeads (ThermoFisher)
- PCR reagents
- Primers
- SPRI Beads (BioDynami Cat.# 40051) or equivalent
- Mineral oil

Protocol

Step 1. 1st Hybridization

1. Warm probe at 37°C, invert the tube several times to dissolve. Mix the following in a 96-well plate:

MHC-II-B Probe	30
NGS Library (500 ng)	50
Total	80 μ l

2. Overlay a drop of mineral oil in wells with above mix. Seal the wells with cap or sealing tape.
3. Heat at 95°C for 5 min
4. Transfer plate to a hybridization oven, hybridize with the following condition:
 - 1) 75°C for 6-8 hrs
 - 2) 70°C overnight
 - 3) 65°C for 6-8 hrs
 - 4) 60°C overnight

Step 2. 1st Capture

Note: Warm the W1 and W2 buffer at 37°C until the buffer is clear. Invert the bottle several times to mix.

1. Resuspend **MyOne Streptavidin C1 Dynabeads**, transfer 20 μ l of Dynabeads to 96-well plate
2. Place plate on magnet for 1-2 min, aspirate supernatant.
3. Remove plate from magnet and resuspend Dynabeads with 14 μ l of **RS Buffer**.
4. Transfer 80 μ l of hybridized sample to 14 μ l of Dynabeads solution, mix by pipette. Incubate at room temperature for 5 min.
5. W1 washing:
 - a. Place plate on magnet for 1-2 min, aspirate supernatant.
 - b. Remove plate from magnet, add 200 μ l of **W1 Buffer**, resuspend Dynabeads by pipetting, and incubate at room temperature for 2 min.
 - c. Place plate on magnet for 1-2 min, and aspirate supernatant.
 - d. Repeat W1 washing one more time.
6. Add 200 μ l of **80% ethanol** gently without disturbing Dynabeads, incubate at room temperature for 30 sec, aspirate supernatant COMPLETELY.
7. Air dry beads for 4 min.
8. Remove plate from magnet and resuspend Dynabeads in 26 μ l of water. Seal the plate and heated at 95°C for 3 min, chill plate on ice.
9. Place plate on magnet, transfer 23 μ l of supernatant to a new PCR plate

Step 3. PCR: Amplify captured NGS library. Use below reaction as a reference.

1. Mix the following in 96-well PCR plate:

Captured library (from step 2)	23	
Primers*	2	* Not included in the kit
2X PCR mix*	25	
Total	50 μ l	

2. Put PCR plate on a thermal cycler, start PCR with the following condition:

Step	Temperature	Time	Cycles
Initial Denaturation	98°C	30 seconds	1
Denaturation	98°C	10 seconds	10 cycles
Annealing/extension	65°C	70 seconds	
Final Extension	65°C	2 minutes	1
Hold	4°C		

Step 4. 2nd Hybridization

1. Warm probe at 37°C, invert the tube several times to dissolve. Mix the following in a 96-well plate:

MHC-II-B Probe	30
PCR mixture (from step 3)	50
Total	80 µl

2. Overlay a drop of mineral oil in wells with above mix. Seal the wells with cap or sealing tape.
3. Incubate at 95°C for 5 min
4. Put plate in the hybridization oven, hybridize with the following condition:
 - 1) 75°C for 6-8 hrs
 - 2) 70°C overnight
 - 3) 65°C for 6-8 hrs
 - 4) 60°C overnight

Step 5. 2nd Capture

Note: Warm the W1 and W2 buffer at 37°C until the buffer is clear. Invert the bottle several times to mix.

1. Resuspend **MyOne Streptavidin C1 Dynabeads**, transfer 20 µl of Dynabeads to 96-well plate.
2. Place plate on magnet for 1-2 min, aspirate supernatant.
3. Remove plate from magnet and resuspend Dynabeads with 14 µl of **RS Buffer**.
4. Transfer 80 µl of hybridized sample to 14 µl of Dynabeads solution, mix by pipette. Incubate at room temperature for 5 min.
5. W1 washing:
 - a. Place plate on magnet for 1-2 min, aspirate supernatant.
 - b. Remove plate from magnet, add 200 µl of **W1 Buffer**, resuspend Dynabeads by pipetting, and incubate at room temperature for 2 min.
 - c. Place plate on magnet for 1-2 min, and aspirate supernatant.
6. W2 washing:
 - a. Remove plate from magnet, add 200 µl of **W2 Buffer**,
 - b. Resuspend Dynabeads by pipetting, incubate plate at 65°C (water bath, heat block or thermal cycler) for 5 min,
 - c. Place plate on magnet for 10 sec, aspirate supernatant.
 - d. Repeat W2 washing one more time.
7. Add 200 µl of **80% ethanol** gently without disturbing Dynabeads, incubate at room temperature for 30 sec, aspirate supernatant COMPLETELY.
8. Air dry beads for 4 min.
9. Remove plate from magnet and resuspend Dynabeads in 26 µl of water. Seal the plate and heated at 95°C for 3 min, chill plate on ice.
10. Place plate on magnet, transfer 23 µl of supernatant to a new PCR plate.

Step 6. Final PCR: Amplify captured libraries. Use below reaction as a reference.

1. Mix the following in 96-well PCR plate:

Captured library	23
Primers*	2
2X PCR mix*	25
Total	50 µl

2. Put PCR plate on a thermal cycler, start PCR with the following condition:

Step	Temperature	Time	Cycles
Initial Denaturation	98°C	30 seconds	1
Denaturation	98°C	10 seconds	10-20
Annealing/extension	65°C	70 seconds	cycles**
Final Extension	65°C	2 minutes	1
Hold	4°C		

* Not included in the kit

** As a reference:

10-15 cycles for 4 MB target region;
 11-16 cycles for 2 MB target region;
 12-17 cycles for 1 MB target region;
 13-18 cycles for 500 KB target region;
 14-19 cycles for 250 KB target region;

3. Place PCR plate on ice. Load 5 μ l PCR product on 2% agarose gel for confirmation of amplification. Add more PCR cycles if needed.

Step 7. Beads Purification

1. Resuspend **SPRI beads** and transfer 40 μ l to the above reaction mixture, mix by pipetting and incubate for 3 min.
2. Load the plate on a magnet, incubate for 2 min, and discard the supernatant carefully.
3. Add 180 μ l of **80% ethanol**, incubate for 30 sec, and discard the supernatant carefully. Remove all residual ethanol without disturbing the beads.
4. Remove the plate from the magnet, resuspend the beads in 22 μ l of **water** or **Tris-HCl** (10 mM).
5. Load the plate on the magnet, incubate for 1 min, and transfer 20 μ l supernatant (containing library) to a new tube without disturbing the beads.

Quality Control

Kit components passed stringent functional quality test.

Product Use Limitation




This product is developed and sold for research purposes and *in vitro* use only. Please refer to BioDynamy.com for Material Safety Data Sheet of the product.

Limited Label License

The product is developed and sold exclusively for research purposes and *in vitro* use only. The product or its any individual component has not been tested for use in diagnostics or drug development, and is not suitable for administration to human or animal.

The purchaser of this product is granted a limited, non-transferable right to use the purchased amount of the product only for internal, research purposes for the sole benefit of the purchaser. The buyer cannot sell or otherwise transfer (i) this product (ii) its components or (iii) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for commercial purposes. This product is for internal research purposes only and is not for use in commercial purposes of any kind. "Commercial purposes" includes any activity for which a party receives consideration and may include, but is not limited to, (1) use of the product or its components or derivatives in manufacturing, (2) transfer or sale of vectors made with the product or components or derivatives of the product, (3) use of this product or components or derivatives of the product made therefrom to provide a service, information, or data to a third party in return for a fee or other consideration, or (4) resale of the product or its components or derivatives, whether or not such product or its components or derivatives are resold for use in research. If the purchaser is not willing to accept the limitations of this limited use statement, BioDynamy is willing to accept return of the products with a full refund. For information on obtaining additional rights, please contact support@biodynamy.com

BioDynamy

-  601 Genome Way, Huntsville, Alabama 35806, USA
-  <https://biodynamy.com>
-  support@biodynamy.com



Dec. 2022